



Waisman Center
Motor Speech Conference
Savannah, Georgia
4 March 2010

Muscle-fiber heterogeneity in craniofacial muscles: Implications for speech development and speech motor control

Ray D. Kent, Ph.D.


Waisman Center

University of Wisconsin - Madison



INTRODUCTION:

The essential enzyme of muscle contraction is myosin, which hydrolyzes ATP to provide the energy that activates actin-myosin crossbridges to generate force or to do physical work. The various subunits of myosin are expressed as families of isoforms that have the same general function but allow for variations in quantitative aspects such as contraction rate and metabolism. The different muscle fiber types create a continuum of different contraction speeds ranging from low to high as follows.



The continuum is enhanced by the presence of developmental isoforms (e.g., Fetal), specialized isoforms (e.g., Mandibular, Cardiac), and hybrid fibers (e.g., IM/IIC). Hybrid fibers usually have a contraction speed that is intermediate to their constituent pure isoforms.

METHODS:

A review of the literature was conducted using High Wire Press, PubMed, and other literature retrieval systems. Search terms included: muscle fibers, craniofacial muscles, myosin heavy chain, and the names of individual muscles. The primary goal was to collate information on the relative proportion of muscle-fiber types, but information on other properties, such as metabolism and developmental transitions, was also noted. Muscle fiber data were obtained for the following muscle groups, with individual muscles identified in parentheses:

- Facial** (orbicularis oris, buccinator, zygomatic)
- Lingual** (genioglossus; superior, longitudinal, and transverse intrinsic muscles)
- Jaw elevators** (masseter, medial pterygoid, temporalis)
- Jaw depressors** (digastric, geniohyoid, mylohyoid)
- Palatal elevators** (levator veli palatini, tensor veli palatini)
- Palatal depressors** (palato-pharyngeus, uvula)
- Pharyngeal** (cricopharyngeus, superior pharyngeal constrictor, cricothyro-pharyngeus)
- Laryngeal** (posterior cricoarytenoid, cricothyroid, lateral cricoarytenoid, thyroarytenoid, vocalis).

RESULTS:

Main results are summarized with arithmetic signs:
= (roughly equal), > (more than) and >> (much more than) to show the relative number of different fiber types in a muscle. For example, a muscle with a predominance of type I fibers and only a few type IIA fibers would be expressed as I >> IIA.

Facial muscles

Orbicularis oris (OO): IIA >> I = IIC
Zygomatic (Zyg): I = IIA > IIC

Implications:

Abundance of type II fibers in OO is consistent with the rapid acceleration and high speed needed for intermittent orofacial movements, such as bilabial stops. NB: OO and Zyg share embryologic origin and innervation but differ in function; therefore muscle fiber composition seems to be determined largely by function.

Tongue muscles

Intrinsic muscles in anterior tongue: IIA >>> I = IM/IIC
Intrinsic muscles in posterior tongue: I = IM/IIC > IIA

Implications:

The predominance of small type IIA fibers in the anterior portion permits rapid movements of tongue tip and blade, whereas a proportionately larger population of larger type I and type IM/IIC fibers in the posterior region aids postural support of the tongue body and relatively slow vowel-to-vowel movements.

Jaw muscles

Jaw elevators: I = Hybrid >> IIA = IIX = Fetal = Cardiac
Suprahyoid depressors: I = IIA > IIX > Hybrid
Infrahyoid depressors: I = IIA >> IIX = Hybrid

Implications:

The jaw elevators are adapted for relatively slow, tonic movements such as postural support for speech and swallowing movements, whereas the depressors are suited to faster, phasic movements, such as jaw opening for syllables, and presumably consonant-to-vowel transitions in babbling. Fiber polymorphism enables the jaw muscles to achieve different specializations in chewing, swallowing, and speech.

Pharyngeal muscles

Pharyngeal constrictor: IIA > I > Hybrid
Cricopharyngeal: IIA > I > Hybrid
Cricothyropharyngeus: I > IIA = Tonic > IIX
Other histological features

Implications:

Studies document substantial inter-individual variability, presence of hybrid fibers, and a predominance of Type IIA fibers. Mu and Sanders (2008), in their description of a newly discovered muscle, the cricothyropharyngeus, noted unusual MyCH isoforms including slow-tonic, alpha-cardiac, neonatal, and embryonic. They concluded that this muscle "appears to be a newly described and uniquely human muscle with characteristics suggesting a specialized function that may be speech related" (p. 927).

Palatal muscles

Palate elevators: I >> IIA = IIAAB
Palate depressors: IIA = IIAAB > I

Implications:

Elevators are designed for tonic contraction to support prolonged velopharyngeal closure as needed for non-nasal sound sequences (e.g., canonical babbling), whereas depressors are more suited to rapid phasic movements of the palate. The agonist-antagonist muscles have a different fiber composition that appears to meet functional needs.

Laryngeal muscles

Posterior cricoarytenoid (PCA): I >> IIA = Hybrid
Lateral cricoarytenoid (LCA): I = IIA = IIX
Thyroarytenoid (TA): I = IIX > IIA > Hybrid
Vocalis: I = IIA = IIX
Cricothyroid: IIA > I = Tonic

Implications:

The PCA, the laryngeal abductor, has a slower fiber type profile than the principal adductor, the TA, a difference that may be related to the tonic activity of the PCA during respiration.

Development of the vocal fold can be summarized as:

- 1) monolayer of cells in the neonate
- 2) bilaminar structure by about two months
- 3) a more fully constituted bilaminar structure by five months
- 4) a trilayer by the age of seven years

Therefore, the lamina propria undergoes substantial change during early periods of vocal experience. A study of unphonated vocal folds in three young adults with severe cerebral palsy evinced abnormalities in vocal fold mucosa presumably due to the lack of mechanical stimulation normally provided by phonation (Sato, Nakashima, Nonaka, & Harabuchi, 2008).

SUMMARY:

- a) Slower-type profiles are found in: zygomatic, jaw elevators, posterior tongue, palatal elevators, and the laryngeal abductor (posterior cricoarytenoid). These profiles are suited to tonic contractions and postural support.
- b) Faster-type profiles are found in: lips (orbicularis oris), jaw depressors, anterior tongue, palatal depressors, pharyngeal muscles, and laryngeal adductors. These profiles are suited to rapid phasic movements.

GENERAL PROPERTIES OF MUSCLE FIBERS IN CRANIOFACIAL MUSCLES:

1. These muscles are unique in their genetic, developmental, functional, and phenotypical properties (Schuler & Dalrymple, 2001; Tzahor, 2009).
2. The muscle-fiber composition of these muscles differs (a) from muscles of the limbs or trunk, (b) from homologous muscles in nonhuman species, (c) across muscles and sometimes from one region or belly of a muscle to another, and (d) across agonist-antagonist muscle sets (e.g., laryngeal adductors/abductors, jaw depressors/elevators, velar depressors/elevators) (reviewed in Kent, 2004).
3. The degree of hybridism in muscle fibers may reflect the diversity or specialization of activity in a given muscle (Hoh, 2002, 2005; Wu, Crumley, & Caiozzo, 2000; Stephenson, 2006). Speech is a beneficiary of this feature.
4. Variation in muscle-fiber types within a muscle, together with localized motor unit territories, enables task-specific motor activity (Van Eijden & Turkawski, 2001).
5. Speech motor control is a beneficiary of these properties, which have general relevance in explaining certain aspects of speech development including motor patterns in babbling.

REFERENCES:

- A full set of references is available at <http://www.waisman.wisc.edu/vocal/>.
- Kent, R. D. (2004). The uniqueness of speech among motor systems. *Clinical Linguistics & Phonetics*, 18, 495-505.
- Hoh, J. F. Y. (2002). 'Superfast' or masticatory myosin and the evolution of jaw-closing muscles of vertebrates. *Journal of Experimental Biology*, 205, 2203-2210.
- Hoh, J. F. Y. (2005). Laryngeal muscle fibre types. *Acta Physiologica Scandinavica*, 163, 13-149.
- Mu, L., & Sanders, I. (2008). Newly revealed cricothyropharyngeus muscle in the human laryngopharynx. *Anatomic Record*, 291, 927-938.
- Sato, K., Nakashima, T., Nonaka, S., & Harabuchi, Y. (2008). Histopathologic investigations of the unphonated human vocal fold mucosa. *Acta Otolaryngologica*, 128, 694-701.
- Schuler, C. R., & Dalrymple, K. R. (2001). Molecular regulation of tongue and craniofacial muscle differentiation. *Critical Reviews in Oral Biology and Medicine*, 12, 3-17.
- Stephenson, G. M. M. (2006). Cognitive value of hybrid fibres. *Brazilian Journal of Morphology*, 23, 187-194.
- Tzahor, E. (2009). Heart and craniofacial muscle development: a new developmental theme of distinct myogenic fields. *Developmental Biology*, January 6, 2009.
- Van Eijden, T. M., & Turkawski, S. J. (2001). Morphology and physiology of masticatory muscle motor units. *Critical Reviews in Oral Biology & Medicine*, 12, 76-91.
- Wu, Y. Z., Crumley, R. L., & Caiozzo, V. J. (2000). Are hybrid fibers a common motif of canine laryngeal muscles? *Archives of Otolaryngology-Head & Neck Surgery*, 126, 865-873.

ACKNOWLEDGMENTS:

This work was supported by NIH/NIDCD grants # R03-DC 4362 & R01-DC 006282 to the Vocal Tract Development Lab. Also, a core grant to the Waisman Center from NIH-NICHD P30-HD03352. Thanks to Jennifer J. Lewandowski for her assistance in poster preparation.

Bibliography for Craniofacial Muscle Fibers in Humans

- Boseley, M. E., & Hartnick, C. J. (2006). Development of the human true vocal fold: depth of cell layers and quantifying cell types within the lamina propria. *Annals of Otolaryngology, Rhinology and Laryngology*, 115, 784-788.
- Bottinelli, R. (2001). Functional heterogeneity of mammalian single muscle fibres: do myosin isoforms tell the whole story? *Pflügers Archive*, 443, 6-17.
- Grunheid, T., Langenbach, G. E. J., Korfage, J. A. M., Zentner, A., & van Eijden, T. M. G. J. (2009). The adaptive response of jaw muscles to varying functional demands. *European Journal of Orthodontics*, doi: 10.1093/ejo/cjp093.
- Han, Y., Wang, J., Fischman, D. A., Biller, H. F., & Sander, I. (1999) Slow tonic muscle fibers in the thyroarytenoid muscles of human vocal folds: a possible specialization for speech. *Anatomic Record*, 256, 146-157.
- Hannam, A. G., & McMillan, A. S. (1994). Internal organization in the human jaw muscles. *Critical Reviews in Oral Biology and Medicine*, 5, 55-89.
- Hartnick, C. J., Rehbar, R., & Prasad, V. (2005). Development and maturation of the pediatric human vocal fold lamina propria. *Laryngoscope*, 115, 4-15.
- Hirano, M., Kurita, S., & Hakashima, T. (1983). Growth, development, and aging of human vocal folds. *Vocal fold physiology: contemporary research and clinical issues* (pp. 23-43).
- Hoh, J. F. Y. (2002). 'Superfast' or masticatory myosin and the evolution of jaw-closing muscles of vertebrates. *Journal of Experimental Biology*, 205, 2203-2210.
- Hoh, J. F. Y. (2005). Laryngeal muscle fibre types. *Acta Physiologica Scandinavica*, 183, 13-149.
- Hwang, K., Kim, D. J., & Hwang, S. H. (2007). Immunohistochemical study of differences between the muscle fiber types in the pars peripheralis and marginalis. *Journal of Craniofacial Surgery*, 18, 591-593.
- Kent, R. D. (2004). The uniqueness of speech among motor systems. *Clinical Linguistics & Phonetics*, 18, 495-505.
- Korfage, J. A., Brugman, P., & Van Euden, T. M. (2000). Intermuscular and intramuscular differences in myosin heavy chain composition of the human masticatory muscles. *Journal of Neurological Science*, 178, 95-106.
- Korfage, J. A. M., Hilmers, R., de Gouyon Matignon, M., van Wessel, T., Langenbach, G. E. J., & van Eijden, T. M. G. J. (2009). Postnatal development of

fiber type composition in rabbit jaw and leg muscles. *Cells, Tissues, Organs*, 190, 42-52.

Korphage, J. A. M., & Van Eigden, T. M. G. J. (2003). Myosin heavy chain composition in human masticatory muscles by immunohistochemistry and gel electrophoresis. *Journal of Histochemistry & Cytochemistry*, 51, 113-119.

Li, Z. B., Lehar, M., Nakagawa, H., Hoh, J. F., & Flint, P. W. (2004). Differential expression of myosin heavy chain isoforms between abductor and adductor muscles in the human larynx. *Otolaryngology--Head and Neck Surgery*, 130, 217-222.

Lloyd, J. S., Brozanski, B. S., Daood, M., & Watchko, J. F. (1996). Developmental transitions in the myosin heavy chain phenotype of human respiratory muscle. *Biology of the Neonate*, 69, 67-75.

McLoon, L. K., Thorstenson, K. M., Solomon, A., & Lewis, M. P. (2007). Myogenic precursor cells in craniofacial muscles. *Oral Diseases*, 13, 134-140.

Mu, L., & Sanders, I. (2008). Newly revealed cricothyropharyngeus muscle in the human laryngopharynx. *Anatomic Record*, 291, 927-938.

Mu, L., Wang, J., Su, H., & Sanders, I. (2007). Adult human upper esophageal sphincter contains specialized muscle fibers expressing unusual myosin heavy chain isoforms. *Journal of Histochemistry & Cytochemistry*, 55, 199-207.

Pae, E. K., Hyatt, J. P., Wu, J., & Chien, P. (2007). Short-term electrical stimulation alters tongue muscle fibre type composition. *Archives of Oral Biology*, 52, 544-551.

Perie, S., Agbulut, O., St. Guily, J. L., & Butler-Brown, G. S. (2000). Myosin heavy chain expression in human laryngeal muscle fibers. A biochemical study. *Annals of Otolaryngology, Rhinology, and Laryngology*, 109, 216-220.

Perie, S., Lacau St. Guily, J., Callard, P., & Sebille, A. (1997). Innervation of adult human laryngeal muscles. *Journal of the Neurological Sciences*, 149, 81-86.

Perie, S., St. Gully, J. L., & Sebille, A. (1999). Comparison of perinatal and adult multi-innervation in human laryngeal muscles. *Annals of Otolaryngology, Rhinology and Laryngology*, 108, 683-688.

Porter, J. D., Israel, S., Gong, B., Merriam, A. P., Feurerman, J., Khanna, S., & Kaminski, H. J. (2006). Distinctive morphological and gene/protein expression signatures during myogenesis in novel cell lines from extraocular and hindlimb muscle. *Physiological Genomics*, 24, 264-275.

Reiser, P. J. (in press). Myosin expression patterns in dog thyroarytenoid muscle: insights into muscle function from single fibers. *Communication Disorders Review*.

Ren, M., & Mu, L. (2005). Intrinsic properties of the adult human mylohyoid muscle: neural organization, fiber-type, and myosin heavy chain expression. *Dysphagia*, 20, 182-194.

Rosenberg, T. L., & Schweinfurth, J. M. (2009). Cell density of the lamina propria of neonatal vocal folds. *Annals of Otology, Rhinology & Laryngology*, 118, 87-90.

Roy, R. R., Pierotti, D. J., Garfinkel, A., Zhong, H., Baldwin, K. M., & Edgerton, V. R. (2008). Persistence of motor unit and muscle fiber types in the presence of inactivity. *Journal of Experimental Biology*, 211, 1041-1049.

Saigusa, H., Niimi, S., Yamashita, K., Gotoh, T., & Kumada, M. (2001). Morphological and histochemical studies of the genioglossus muscle. *Annals of Otology, Rhinology & Laryngology*, 110, 779-784.

Sato, K., & Hirano, M. (1995). Histologic investigation of macula flava of the human newborn vocal fold. *Annals of Otology, Rhinology and Laryngology*, 104, 556-562.

Sato, K., Nakashima, T., Nonaka, S., & Harabuchi, Y. (2008). Histopathologic investigations of the unphonated human vocal fold mucosa. *Acta Otolaryngologica*, 128, 694-701.

Sato, M., & Sato, T. (1992). Fine structure of developed human tongue muscles. *Okajimas Folia Anatomica Japan*, 69, 115-130.

Schiaffino, S., Reggiani, C. (1996). Molecular diversity of myofibrillar proteins: gene regulation and functional significance. *Physiological Review*, 76, 371-423.

Schuler, C. R., & Dalrymple, K. R. (2001). Molecular regulation of tongue and craniofacial muscle differentiation. *Critical Reviews in Oral Biology and Medicine*, 12, 3-17.

Sciote, J. J., Horton, M. J., Rowlerson, A. M., & Link, J. (2003). Specialized cranial muscles: How different are they from limb and abdominal muscles? *Cells, Tissues, Organs*, 174, 73-86.

Sciote, J. J., Morris, T. J., Brandon, C. A., Horton, M. J., & Rosen, C. (2002). Unloaded shortening velocity and myosin heavy chain variations in human laryngeal muscle fibers. *Annals of Otology, Rhinology, & Laryngology*, 111, 120-127.

- Sciote, J. J., Rowleron, A. M., Hopper, C., & Hunt, N. P. (1994). Fibre type classification and myosin isoforms in the human masseter muscle. *Journal of Neuroscience*, 126, 15-24.
- Scott, W., Stevens, J., & Binder-Macleod, S. A. (2001). Human skeletal muscle fiber type classifications. *Physical Therapy*, 81, 1810-1816.
- Sinanan, A. C., Hunt, N. P., & Lewis, M. P. (2004). Human adult craniofacial muscle-derived cells: neural-cell adhesion-molecule (NCAM; CD56)-expressing cells appear to contain multipotential stem cells. *Biotechnology and Applied Biochemistry*, 40, 25-34.
- Slaughter, K., Li, H., & Sokoloff, A. J. (2005). Neuromuscular organization of the superior longitudinalis muscle in the human tongue. 1. Motor endplate morphology and muscle fiber architecture. *Cells, Tissues, Organs*, 181, 51-64.
- Smith, J. C., Goldberg, S. J., & Shall, M. S. (2005). Phenotype and contractile properties of mammalian tongue muscles innervated by the hypoglossal nerve. *Respiratory Physiology & Neurobiology*, 147, 253-262.
- Stal, P. (1994). Characterization of human oro-facial and masticatory muscles with respect to fibre types, myosins and capillaries. Morphological, enzyme-histochemical, immune-histochemical and biochemical investigations. *Swedish Dental Journal* [Supplement], 98, 1-55.
- Stal, P., Eriksson, P. O., Eriksson, A., & Thornell, L. E. (1990). Enzyme-histochemical and morphological characteristics of muscle fibre types in the human buccinators and orbicularis oris. *Archives of Oral Biology*, 35, 449-458.
- Stal, P., & Lindman, R. (2000). Characterisation of human soft palate muscles with respect to fibre types, myosins and capillary supply. *Journal of Anatomy*, 197, 275-290.
- Stal, P., Marklund, S., Thornell, L. E., DePaul, R., & Eriksson, P. O. (2003). Fibre composition of human intrinsic tongue muscles. *Cells, Tissues, Organs*, 173, 147-161.
- Stephenson, G. M. M. (2006). Cognitive value of hybrid fibres. *Brazilian Journal of Morphology*, 23, 187-194.
- Sundman, E., Ansved, T., Margolin, G., Kuylenstierna, R., & Eriksson, L. I. (2004). Fiber-type composition and fiber size of the human cricopharyngeal muscle and the pharyngeal constrictor muscle. *Acta Anaesthesiologica Scandinavica*, 48, 423-429.
- Surchak, A., Hunt, N. P., Shah, R., Sinanan, A. C. M., Loyd, T., & Lewis, M. P. (2008). Myosin proteins identified from masseter muscle using quantitative

reverse transcriptase-polymerase chain reaction—a pilot study of the relevance to orthodontics. *European Journal of Orthodontics*, doi:10.1093/ejo/chn095.

Tzahor, E. (2009). Heart and craniofacial muscle development: a new developmental theme of distinct myogenic fields. *Developmental Biology*, January 6, 2009.

Van Eijden, T. M., & Turkawski, S. J. (2001). Morphology and physiology of masticatory muscle motor units. *Critical Reviews in Oral Biology & Medicine*, 12, 76-91.

Wang, J., Han, Y., Su, H., & Mu, L. (2004). Expression of unique and developmental myosin heavy chain isoforms in adult human digastric muscle. *Journal of Histochemistry and Cytochemistry*, 52, 851-859.

Wu, Y. Z., Crumley, R. L., Armstrong, W. B., & Caiozzo, V. J. (2000). New perspectives about human laryngeal muscles: single-fiber analyses and interspecies comparisons. *Archives of Otolaryngology-Head & Neck Surgery*, 126, 837-864.

Wu, Y. Z., Crumley, R. L., & Caiozzo, V. J. (2000). Are hybrid fibers a common motif of canine laryngeal muscles? *Archives of Otolaryngology-Head & Neck Surgery*, 126, 865-873.

Yu, F., Stal, P., Thornell, L. E., & Larson, L. (2002). Human single masseter muscle fibers contain unique combinations of myosin and myosin binding protein C isoforms. *Journal of Muscle Research and Cell Motility*, 23, 317-326.